

PRE-PROCEDURE IMPORTANT INFORMATION

PLEASE READ AT LEAST FIVE DAYS BEFORE YOUR PROCEDURE

At least two days prior to your procedure, the procedure center will text / call to confirm your arrival time and review information regarding your procedure.

PLEASE BE ADVISED OF THE FOLLOWING:

It is your responsibility to contact your insurance company to learn whether a specific procedure is covered and if it will be applied to a deductible. Some insurance plans have riders and underwriting on the plan that may or may not cover any or all of the procedure. We want you to be aware and understand the specifics of your insurance coverage. We are happy to provide you with the codes that could be used when submitting your claim to the insurance company.

Depending on the location of your procedure, that locations billing department will verify your insurance benefits and inform you prior to your procedure day if a facility payment is required at the time of service. Anesthesia, pathology, and physician charges will be filed with insurance for determination of additional patient financial responsibility.

VERY IMPORTANT

There are three (3) entities involved that will send a bill following your procedure. You can call the phone numbers provided for each location if you have questions regarding procedure estimates or billing.

- 1. FACILITY: Associated Endoscopy, Mid-State Endoscopy, or Summit Medical Center.
- 2. **ANESTHESIA**: Depending on the location of your procedure, you may receive a bill from the anesthesia group affiliated with the surgery center or hospital where you had your procedure.
- 3. OneGI: Provider & Pathology if any specimens are collected during procedure (615-455-5732)

Anesthesia, pathology, and physician charges are filed with insurance for determination of additional patient financial responsibility.

SURGERY CENTER LOCATIONS:

You will report to one of the surgery centers noted below as instructed by your provider/nurse.

Associated Endoscopy

5653 Frist Boulevard, Suite 532

Hermitage, TN 37076

Phone: 855-880-9327

*Located next door to our Hermitage Office in Summit Medical Center.

Mid-State Endoscopy

1115 Dow Street, Suite A Murfreesboro, TN 37130

Phone: 615-848-9234

*Located next door to Mid-State Gastroenterology (Suite B).

Summit Medical Center Same Day Surgery (SDS)

5653 Frist Boulevard Hermitage, TN 37076 SDS Phone: 615-316-3570

Billing: 800-370-1983

*Check-in on 1st floor at Registration located on left when you enter hospital.

Please note Summit Medical Center offers free valet parking at the patient entrance closest to the ER.



On the day of your procedure:

- 1. You will be required to have someone available to drive you home after the procedure. You cannot and will not be allowed to drive following the procedure. You will not be released to any public mode of transportation (i.e. taxi, uber, Lyft). Due to the effects of the medications, you may not remember the instructions given to you after the procedure. Your driver must stay on the premises while you are having the procedure and be available to speak with the physician and/or staff regarding your post-procedure instructions and findings.
- 2. Please leave all your valuables at home except for your photo ID, insurance card, and any copayment you may be required to bring for the procedure. If your insurance has changed since your procedure was scheduled, you must contact Associates in Gastroenterology at 615-885-1093 and provide them with this new information.
- 3. Complete the <u>PATIENT MEDICATION RECONCILIATION FORM</u> provided and bring with you to your procedure appointment. It is very important that the nursing staff and physician know what medications you are taking (including over-the-counter medicines and herbal supplements). Please be sure to include the dosage you are taking and the date of your last dose.
- 4. All aspirin and aspirin type products, including blood thinners, should have been held as instructed by your physician. This includes Plavix, Warfarin, Ticlid and Coumadin. Please contact your AiG nurse at 615-885-1093 with any questions regarding these medications.
- 5. You may not eat or drink anything for at least four (4) hours prior to your procedure unless it is related to your prep instructions. Please follow the necessary bowel preparation instructions provided for your procedure. No mints, gum, cough drops, ice chips, or sips of any liquid for four (4) hours prior to your procedure. No recreational drugs or alcohol for 24 hours prior to your procedure.
- 6. The only medications you should take the day of your procedure should be for blood pressure or heart medications, unless otherwise instructed.
- 7. Wear comfortable clothing. DO NOT wear contacts. Glasses are permitted.



Colonoscopy Prep General Instructions

- The day before your procedure, you must stay on a <u>Clear Liquid Diet</u>.
 - Clear Liquid Diet

IMPORTANT NOTE Do not consume any red-, pink-, or purple-colored liquids or foods.

- Coffee & Tea (no dairy or cream)
- Fruit Juice without Pulp (apple, white grape, white cranberry)
- Gatorade, G2, Powerade
- Crystal Light
- Broth
- Jell-O
- Sodas (Coke, Diet Coke, etc.)
- Juice Popsicles without Fruit or Pulp
- DO NOT consume any milk/cream products, red-, pink-, or purple-colored liquids, or any alcoholic beverages.
- Drink 6-8 ounces of clear liquid every hour throughout the day to stay hydrated.
- Nothing by mouth four (4) hours prior to procedure, including gum, mints, candy, or water.
- Do not use any recreational drugs or alcohol for 24 hours prior to your procedure.

Medications

- You will need to stop taking full-dose aspirin (325mg), blood thinners and/or arthritis medications such as Motrin, Advil, Vitamin E, and herbal supplements for 5 days prior to your procedure, unless otherwise instructed. You may take Baby Aspirin (81mg), Tylenol, multi-vitamins, and all other prescription medications. If you are unsure about a medication, please ask your physician or nurse. Please see attached list of medications to avoid.
- If you take heart, blood pressure or seizure medications you should take them as directed the morning of your procedure with a small amount of water. All other medications should be held until instructions are given after your procedure.
- GLP-1 medications, such as Dulaglutide (Trulicity) (weekly), Exenatide extended release (Bydureon bcise) (weekly), Exenatide (Byetta) (twice daily), Semaglutide (Ozempic, Wegovy) (weekly), Liraglutide (Victoza, Saxenda) (daily), Lixisenatide (Adlyxin) (daily), Semaglutide (Rybelsus) MUST BE STOPPED prior to your procedure and taken as follows:
 - o If taken once a day, hold the medication the day of the procedure.
 - o If taken once a week, hold the medication one (1) week prior to your procedure.
 - o Consult with your endocrinologist if taking GLP-1 agonist for diabetes.

Don't Forget

- You must drink the recommended amount of water and/or clear liquids while taking this bowel preparation to prevent serious side effects and to ensure that you are properly prepared.
- If you do not follow these directions, your exam may have to be repeated or rescheduled.
- You must have a responsible adult to accompany you home. If you do not have an adult escort, your procedure will be cancelled and rescheduled.
- You need to bring your insurance card and a photo ID with you.
- Please bring your medication list with you on the day of your procedure, along with the date your medication was last taken.

PROCEDURE DATE	
PROCEDURE TIME	





PLENVU PREP INSTRUCTIONS

PLEASE BE SURE TO FOLLOW THESE INSTRUCTIONS (DO NOT FOLLOW THE ONES ON THE PACKAGE)

Evening Before Procedure @ 6PM

Use the mixing container to mix the contents of the Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time and slowly finish the dose within 30 minutes. Refill container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.

Day of Colonoscopy

Six (6) hours before your procedure, use the mixing container to mix the contents of Dose 2, pouches A and B, with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time and slowly finish the dose within 30 minutes. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.

After drinking this, nothing else by mouth prior to your procedure.

IMPORTANT:

- IF YOU EAT OR DRINK ANYTHING WITHIN FOUR (4) HOURS OF YOUR PROCEDURE, THIS
 WILL RESULT IN A DELAY OR CANCELLATION OF YOUR PROCEDURE!! THIS INCLUDES
 WATER, MINTS, HARD CANDY, AND GUM!!
- NO USAGE OF MARIJUANA, COCAINE, RECREATIONAL DRUGS, OR ALCOHOL WITHIN 24 HOURS OF YOUR PROCEDURE. Anesthesia will not sedate you if you have consumed any of these in 24 hours.

PROCEDURE DATE	
PROCEDURE TIME	

PRODUCTS TO AVOID PRIOR TO YOUR PROCEDURE

(IMPORTANT: These are drugs that can thin your blood and may cause bleeding.)

Avoid these medications for at least 5 days before your procedure.

Advil	Loritab ASA		
Aggrenox	Magsal		
Aleve	Midol		
Alka Seltzer	Mobic/Meloxicam		
Anaprox	Mobigesic		
Anacin	Monogesic Tablets		
Arthritis Pain Formula	Motrin		
Ascriptin	Naprosyn		
Aspergum	Naprelan		
Aspirin 325mg (Baby Aspirin 81mg is OK)	Naproxen		
Azoid	Norgesic		
BC Powders	Norwich		
Bextra	Nuprin		
Brilinta	P-A-C Analgesic		
Bufferin	Pepto-Bismal		
Carna Arthritis Medication	Percodan		
Celebrex	Phentermine HCL		
Clinoril	Plaquenil		
Coumadin	Plavix		
Darvon Compound / Darvocet	Pradaxa		
Disalcid	Robaxisal		
Doan's Pills	Salflex		
Dristan	Salsalate		
Easpirin	Salsitab		
Ecotrin	Savaysa		
Effervescent Tablets	Sine-Aid		
Effient	Soma Compound		
Eliquis	Stanback		
Empirin	Synalgos-DC		
Equagesic	Talwin		
Etodolac	Therapy Bayer Caplets		
Excedrin	Tolectin		
Feldene	Tolmetin		
Fenoprex	Toradol		
Fiorinal	Trigesic		
Fish Oil	Trilisate		
Formula Caplets	Urisinus-Inlav		
Goody Powders	Vanquish		
Haltran	Vitamin E		
Ibuprofen	Vioxx		
Ibu-Tab	Warfarin		
Indocin	Xarelto		
Lodine	Zontivity		

PATIENT MEDICATION RECONCILIATION Form

PT. STICKER

Height:		Weight:		Date	of Birth:		Age:
Allergies:	□Yes	□ No Known Allergies	Latex Al	lergy No Y	es 🗆 T	esting Performed	for Latex Allergy
Allergy (Drug) Reaction			Allergy (Drug)	Reaction		
CURRENT PI	RESCRIP	ΓΙ VE MEDICATIONS :					
Name of Medication (print please)		Dose	Last Dose Taken/Time	How Often	Continue After Discharge	Stop After Discharge	
		, SUPPLEMENTS & NON-PE				Continue	Ct
Name	or Medic	ation (print please)	Dose	Last Dose Taken/Time	How Often	<u>Continue</u> After Discharge	Stop After Discharge
NEW MEDIC	ATIONS/	DOSAGES YOU SHOULD TA	AKE AFTEF	R DISCHARGE:			
Name of Medication (print please)			Dose		How Often		
Patient/Resn	onsible l	Person Signature:		1		Date:	
Patient/Responsible Person Signature:							
Physician Signature:							